

CARBON DIOXIDE (CO₂) STORAGE AND SEQUESTRATION OF LAND COVER IN THE LEYTE GEOTHERMAL RESERVATION

Rodel D. Lasco¹, Joveno S. Lales¹, Ma. Theresa Amueyo¹, Ina Q. Guillermo¹,
Agnes C. de Jesus¹, Reiner Medrano², Orlando F. Bajar² and Cirilo V. Mendoza²

¹Environmental Forestry Programme (ENFOR), College of Forestry and Natural Resources,
University of the Philippines, College, 4031 Laguna, Philippines

²PNOC Energy Development Corporation, Merritt Road, Fort Bonifacio, Metro Manila 1201, Philippines

ABSTRACT

The study estimated **the existing carbon (C)** stored and the rate of C sequestration by vegetation in the Leyte Geothermal Reservation, Philippines. For the **20,438 ha** watershed, the total C storage is **3.84 Mt C (74.70 Mt CO₂)** while C sequestration based on biomass change was **47.35 Kt C (773.77 Kt CO₂)**. Relative to power plant emission, the C stored in the reserve is equivalent to more than **22 years** of CO₂ emission. Annual C sequestration is 27% of CO₂ emission per year. For the next **25 years**, two scenarios were **projected**. Under Scenario I ("Business as usual"), the forest reserve will be able to store and sequester more than **32 years** of CO₂ emission from the power plants. Under Scenario II (Accelerated **Reforestation**), the reserve can store and sequester about **34 years** of CO₂ emission.

In terms of cost, the power project operator is spending **P 7.22 per t CO₂ (P4.4 or /US\$ 0.12 per tC)** for every year of C storage and sequestration. For 25 years, the total cost is **P 30.40 per tCO₂ (P 111.5 or/US\$ 2.94 per tC)** which is comparable to the cost of C offset in other tropical countries.

1.0 INTRODUCTION

There is considerable interest on the role of **terrestrial** ecosystems in the global carbon cycle. It is estimated that about 60 Gt C is exchanged between terrestrial ecosystems and the atmosphere every year, with a net terrestrial uptake of 0.7 ±1.0 Gt C (Schimell et al., 1996). The world's tropical forests which **cover** 17.6 M km² contain 428 Gt C in vegetation and soils.

Forest ecosystems play an important role in the climate change problem because they can both be sources and sinks of atmospheric CO₂. They can be managed to assimilate CO₂ via photosynthesis and store carbon in biomass and in soil (Watson et al., 2000). Great attention is focused on tropical forestry to offset carbon emission due to its cost effectiveness, high potential rates of carbon uptake, and associated environmental and social benefits (Brown et al., 2000). Tropical forests have the biggest long-term potential to sequester atmospheric carbon (80% of the world's forests total) by protecting forested lands, slowing deforestation, reforestation, and agroforestry (Brown et al., 1996). However, at present, the world's forests are estimated to be a net source of 1.8 Gt carbon per year primarily because of deforestation, harvesting and forest degradation (Watson et al., 2000). At least twenty percent of all atmospheric CO₂ emissions are from tropical deforestation.

In the Philippines, the role of forest land cover in the national C budgets has been quantified by previous studies (Lasco and Pulhin, 2002; Lasco and Pulhin, 2000). Information on C stocks and fate of C sequestration of specific land cover is still lacking. This study addresses this data gap. The PNOC is managing a largely forested watershed as part of its geothermal reserve in Leyte. The World Bank **awarded** the PNOC-EDC Leyte geothermal project a Global Environmental Facility (GEF) grant to conduct a study on the CO₂ absorption capacity of the vegetation cover in the study area.

2.0 OBJECTIVES OF THE STUDY

The main objectives of the study were to (a) estimate **quantitatively** the **carbon (C) stocks**

and the rate of C sequestration of the different vegetative cover in the Philippine National Oil Company - Energy Development Corporation (PNOC-EDC) geothermal reservation in Leyte, and (b) determine whether the CO₂ emissions of the geothermal power plants could be offset by C stored in the vegetative cover and soil. It is envisioned that the forest vegetation in the reserve will serve as carbon sink that will offset the C emission of the geothermal power.

3.0 MATERIALS AND METHODS

The study was conducted from April 1998 to June 1999 in the 20,438 ha watershed area inside the PNOC-EDC geothermal block located in the island of Leyte, Philippines. The newly-constructed geothermal power plants in the site have a current generating capacity of 640 MWe emitting 644,389 tCO₂/yr.

Five vegetative cover were studied: natural forests, forest plantations, bush/shrublands, mixed agriculture farms and grasslands. The following major C pools were measured: a) above-ground biomass, b) below-ground biomass, c) litter, and d) soil. For natural forest stands, ten 10m by 100m transects were laid out in forest areas representing the range of forest conditions in the forest reserve, primarily elevation. Complete enumeration of all trees (≥ 10 cm dbh) was conducted in each transect. For each tree, the following data were obtained: a) species name, dbh (stem diameter at 1.3m above the ground), b) total height, and c) height at first branch. Tree biomass was computed using the following allometric equation (Brown, 1997):

$$Y = \exp(-2.134 + 2.530 \cdot \ln(D)) \quad (1)$$

where: Y = biomass per tree (kg)
D = dbh (cm).

For understory vegetation (saplings and seedlings ≤ 10 cm dbh), three 1m x 1m plots were randomly laid out in each transect. All individual trees and woody species were harvested. Fresh weight of leaves, branches and stems were determined in the field. Oven dry weight of the different plant parts and wood density were determined in the laboratory. Carbon contents of plant tissues were analyzed. Herbaceous vegetation were collected in three 2m by 2m plots randomly scattered in each transect. The fresh weight, oven dry weight and

C contents were determined. C stored in the litter layer on the forest floor was obtained by randomly laying out 50cm by 50cm frame at 5 locations in each transect. Fresh weight, oven dry weight and C content were determined. Root biomass was estimated by digging three 100 x 100 x 50cm pits in each transect. All roots in the pit were collected and weighed. Oven dry weight and C content were determined in the laboratory. Soil samples were collected from the same pits where the roots were taken. Soil organic C (SOC) and bulk density were analyzed in the laboratory. The total C stored by the forest ecosystem was determined by adding the C in the biomass, litter and soil.

The above-ground tree biomass measurements were conducted twice: at the beginning and towards the end of the study to be able to estimate C sequestration per year. It was assumed that litter biomass and soil organic C will remain unchanged during the observation period.

Brushlands are areas with relatively open canopy and shrub species predominate with a few scattered trees. A similar methodology as described for natural forests was used for brushlands. Ten 10 by 10 m plots were established in representative brushland areas in the forest reserve. C storage and sequestration of above ground biomass, below-ground biomass, litter and soil was estimated following the same procedure outlined above.

Three 10 x 10 m plots were laid out in each of the following plantation species established by PNOC in the reserve: *Gmelina arborea*, *Swietenia macrophylla* and *Acacia mangium*. C storage and sequestration were determined following the same procedures as above.

Two swards were identified in the grassland area. Of the two swards, cogon (*Imperata cylindrica*) occupies a relatively large space within the geothermal block. Old and young swards of talahib (*Saccharum spontaneum*) occupy small patches usually found along roadsides. Two 2m x 2m plots were marked and divided into eight 1m x 1m subplots for quarterly sampling. Aboveground (shoot) and underground parts (rhizomes and roots) were collected for the determination of total dry matter, dead shoot dry weight, rhizome-to-total dry matter ratio, and carbon content. Shoot growth rate was determined from a separate sampling plot.

Table 1. Total C storage of the Leyte geothermal reservation

Land use	C density (t/ha)	Area (ha)	Total C Stored (t)
Forests	392.96	4672.00	1835925.94
Shrubs/Brushlands	186.31	5304.00	988213.17
<i>S. macrophylla</i> plantation	192.09	15.00	2881.31
<i>A. Mangium</i> plantation	275.42	122.00	33601.69
<i>G. arborea</i> plantation	294.16	58.00	17061.19
Grasslands	10.79	2988.00	32240.52
Coconut	196.75	3657.00	71 9496.47
Rice	3.11	181 1.00	5635.65
Abaca/Banana	114.72	1811.00	207759.73
Total		20438.00	384281 5.66
Total CO ₂ equivalent			141 03133.47

For wetland rice, three sites were selected for the wet Season but only one site was available during the dry season. Three 2m x 2m plots were marked in each study site for the determination of total dry matter and partitioning, and carbon content.

For banana, two study sites representing slightly contrasting light environments were chosen. A small block of 4 clumps was marked in each site for non-destructive estimation of dry matter. Additional clumps, however, were marked later to replace typhoon-damaged clumps. Total dry weight of each clump was recorded as the sum of the dry weight of individual plants in the clump.

For coconut, two sites were marked for the study. A block of 4 nut-bearing palms and another block of 3 juvenile palms were marked in each study site for non-destructive determination of dry matter and carbon content. Dry weight of the trunk was calculated using the formula adopted from Friend and Cortey (1994).

Root dry weight was not determined. Hence, total plant dry weight (TPDW) herein recorded refers to the sum of the dry weight of aboveground parts.

Ail plant tissue samples were brought to the Analytical Service Laboratory of the International Rice Research Institute (IRRI) in Los Baños, Laguna where duplicate laboratory analyses for carbon were carried out.

The total costs incurred by PNOC in protecting, maintenance and rehabilitation of the forest reserve was determined. The total cost was divided by the total C stored in the forest reserve to arrive at the cost per C sequestered.

4.0 RESULTS AND DISCUSSION

4.1 Carbon Storage of Various Land Cover

Table 1 presents the results of C analysis. Total mean C storage in the natural forests of the study area amounts to 393 t/ha (Table 1). These are almost equally divided into the biomass (51%) and the soil (49%). The findings are in agreement with those reported in literature where the soil has been found to store at least 30% of total forest C or as much as the biomass (Muora-Costa, 1996 and Lugo and Brown, 1993). Soil organic carbon (SOC) is important because it has the longest residence time among the organic C pools in the forest (Lugo and Brown, 1993 and IGBP Terrestrial Carbon Working Group, 1998). The soil values of this study is reasonable considering that forest soils up to 1 m depth in Venezuela, Papua New Guinea and Colombia could have up to 300 tC/ha (Lugo and Brown, 1993). The total C storage obtained in this study is much higher than earlier estimates for tropical forests in Mt. Makiling, Luzon island (255 tC/ha) and Mindanao island (262 tC/ha) (Lasco et al., 2002; Lasco and Pulhin, 2000). The Mindanao data however did not include SOC.

Table 2. Total C sequestration based on biomass change of the Leyte geothermal reserve

Land use	C Sequestration (t/ha)	Area (ha)	Total C Sequestration (t/yr)
Forests	0.92	4672.00	4298.24
Shrubs/Brushlands	4.29	5304.00	22754.16
<i>S. macrophylla</i> plantation	3.28	15.00	49.20
<i>A. mangium</i> plantation	18.77	122.00	2289.94
<i>G. arborea</i> plantation	8.21	58.00	476.18
Grasslands	0.00	2988.00	0.00
Coconut	4.78	3657.00	17480.46
Rice	0.00	1811.00	0.00
Abaca/Banana	0.00	1811.00	0.00
Total		20438.00	47348.18
Total CO ₂ equivalent			173767.82

As shown in Table 1, three plantation species were monitored in this study: *Sweitenia macrophylla*, *Gmelina arborea*, and *Acacia mangium*. Tree plantations store 192-294 tC/ha (mean = 254 tC/ha) with *A. mangium* having the highest C storage. This is higher than reported values for tree plantations in Mindanao with a mean of 124 tC/ha (Lasco and Pulhin, 2000) because the latter does not include SOC.

C density of brushlands is equal to 186 tC/ha (Table 1). Of this amount, a greater portion (82.7%) is found in the soil. This could be attributed to the lower vegetative cover compared to forested lands.

Undisturbed *I. cylindrica* swards store about 4.7 - 12.1 tC/ha with a mean of 8.5 tC/ha, 60% of which was stored in the aboveground tissues. Accidental fire or deliberate burning will completely release the carbon stored in the tissues back to the atmosphere. The remaining 40% was stored in the modified stems (rhizomes), a fraction of which might have been metabolized *in situ* or re-translocated to the growth points of emerging young shoots.

The carbon storage capacity of *S. spontaneum* was greater than that of *I. cylindrica*. Undisturbed swards stored 7.7 - 20.9 t C/ha (mean = 13.1). Similar to imperata, the shoots stored greater amount (87.5%) of carbon than the roots. Hence, *S. spontaneum* has more C per unit ground area to release back to the atmosphere in case of accidental or deliberate burning.

The average grasslands have a C density of 10.8 t/ha. The results of the study is consistent with the IPCC (Intergovernmental Panel on Climate Change) default value of 10 t/ha of dry matter for crop lands and pasture lands (Houghton, 1997).

The C storage capacity of wetland rice in the project area ranges from 1.9 to 4.4 t C/ha/crop cycle or an average of 3.1 t C/ha/crop cycle. The C storage capacity of banana 114.72 tC/ha (Table 1). While the carbon storage capacity of nut-bearing coconut palms is 197 tC/ha.

The total C storage of the geothermal reserve is calculated to be 3.84 MtC which is equivalent to 14.10 Mt of CO₂ (Table 1). This is about 22 times the annual CO₂ emission from the 653.5 MWe geothermal power plants. Thus, by conserving the C stored in the reserve, the PNOC is preventing the release of CO₂ equivalent to about 22 years operation of the power plants.

4.2 Carbon Sequestration Potential of Various Land Cover

Table 2 presents the C sequestration of plants in the geothermal reserve. Based on biomass accumulation, the C sequestration rate of natural forests is equal to 0.92 tC/ha/yr (Table 2). This is a little lower than the earlier estimate of 1.5 t/ha/yr for Philippine forests (Lasco and Pulhin, 2000).

As expected, fast-growing tree plantations accumulated C faster than natural forests. Mahogany plantations have a C sequestration rate of 3.28 t/ha/yr (Table 2). Mangium plantations accumulate C at the rate of 18.77 t/ha/yr which is much higher than the other plantations species. Gmelina plantations have a C sequestration rate of 15.45 t/ha/yr. Overall, tree plantations have an average sequestration of 10.01 tC/ha/yr.

Brushland areas have a C sequestration rate of 4.29 t/ha/yr. It has been conservatively assumed that grasslands have a net zero sequestration because of regular burning that prevents natural succession from proceeding. However, the potential of grassland areas lies in transforming them to either of the following: natural forest through natural succession, tree plantation or agroforestry farm.

Among all the agricultural farms, only coconut actively sequesters C being a woody perennial. For the annual and non-woody crops such as rice and banana no sequestration is expected because their biomass remain more or less constant.

For the whole study area, the total C sequestration rate based on biomass change is equal to 47 MtC/yr which is equivalent to 174 MtCO₂/yr. This is already 27% of total annual CO₂ emission (644,389 t) of the geothermal power plants. This implies that to sequester all of the emission in one year, an area 3.7 times the same reserve is needed, i.e., about 75,696 ha. Since the PNOC is maintaining more than 100,000 ha area of geothermal reserve in Leyte, this area is likely to be more than sufficient to sequester the annual C emission of the power plants.

The determination of C sequestration rate was constrained by the duration of the study. Ideally, biomass change should be observed for a number of years to increase accuracy. For the same reason, change in SOC was not monitored since this requires a long period of time. However, it is expected that soils in areas that are recovering from disturbance such as reforestation/tree plantations and brushlands are accumulating SOC. This could range from 0.3 to 2 tC/ha/yr in the tropics (Lugo and Brown, 1993). If this is true for the Leyte study area, then the estimated C sequestration rate is still very conservative.

4.3 Future C Storage and Sequestration Rate

Scenario I: "Business as Usual"

This scenario assumes that the reserve is being managed based on present trends and management input, i.e. the various land uses and their respective land area is to be maintained throughout the 25-year life of the power plants as well as their C sequestration rates. This scenario also takes into account the PNOC-EDC plan to reforest 200 ha per year of grasslands and brushlands. Under this scenario, the forested areas, brushlands and coconut plantations will account for most of the increase in annual sequestration rates.

At the end of 25 years, total C storage will be 5.61 Mt C equivalent to 20.6 Mt CO₂. Thus, the amount of C stored in the biomass 25 years hence will be equal to 32 years of plant operation. This implies that by simply letting natural process takes its course through adequate forest protection supported by a modest reforestation program, PNOC-EDC could conserve more than enough C to offset all of its C emission from the power plants.

Scenario II: Accelerated Reforestation

An alternative scenario is to accelerate the reforestation activities in grassland and brushland areas so that all of them are under tree plantation cover within about 10 years. The other landuses are assumed to remain constant. Under such a scenario, the study area will be able to increase C storage to 6.02 MtC at the end of the 25-year period equivalent to 22.10 tCO₂. This is comparable to exactly 34 years of plant operation.

Many other scenarios could be envisioned involving various levels of management input. The purpose of the above is simply to illustrate how C storage will change over time under a given scenario.

4.4 Cost of Carbon Storage and Sequestration

At present, the total forest management cost of the reserve is P17,601,400 per year. In one year's time, the total C in the entire reserve is equal to the sum of C stored at the beginning plus the total C sequestered for the year. For

1999, this amounts to 3.94 Mt C (or 14.47 Mt CO₂). Thus, the cost of a unit of CO₂ stored and sequestered in one year is P1.22 per t CO₂ (P 4.46 per tC). Assuming the same cost of maintenance, it will cost P 30.40 to keep a ton of CO₂ for 25 years (P 111.58/ or S\$ 2.94 per tC).

This unit cost is relatively cheap compared to Thailand where a teak plantation costs US\$ 13-26 per tC (Brown et al., 1996). However, in most other tropical countries the cost of C offset projects is estimated to be US\$ 2-4 per tC (Dixon et al., 1993) which is consistent with the results of the study.

5.0 CONCLUSIONS AND RECOMMENDATIONS

The results show that while the study area cannot absorb all the annual CO₂ emission of the power plants, the amount of C stored in the biomass of the reserve as well as the C to be sequestered for 25 years will more than offset the total C that the power plants will emit in 25 years. Thus, the PNOC-EDC operation in Leyte will not exacerbate the problem of global warming.

To sustain the stored C and the sequestration capacity of the geothermal reservation vegetation, it is important for the project operator to: a) pursue protection of existing natural forests and plantations, b) to manage reservation settlers in place and provide them livelihoods to prevent encroachment of forests, c) to reforest brushlands and grasslands, and d) to encourage the revegetation of built-up areas.

Plant species of higher C sequestration indices can be screened for each land use so that those with greater capacity to absorb C can be prioritized in planting activities.

A land use master plan consulted with the stakeholders with bias towards vegetation establishment that can serve as carbon sink can be formulated.

An inventory of the natural and anthropogenic sources of carbon dioxide inclusive of geothermal power plants can be conducted by government to determine the total C sink requirement so that responsible parties can be assigned to revegetate areas equivalent to their emissions. Thus, while PNOC-EDC as

administrator of the reservation shall maintain the stored C in existing natural forests, the incremental C sink required for the geothermal power plants and other industries can come from reforestation quota of the project proponent.

ACKNOWLEDGMENTS

The authors would like to express its gratitude to the Global Environmental Facility through the World Bank for providing the funds for the conduct of the study: However, the authors bear sole responsibility for the accuracy of the results.

REFERENCES

- Brown, S. (1997). Estimating biomass and biomass change of tropical forest: A primer. Forestry paper 134, FAO. Rome.
- Brown, S., Sathaye, J., Cannel, M., and Kauppi, P. (1996). Management of forests for mitigation of greenhouse gas emissions. Chapter 24 in R.T. Watson, M.C. Zinyowera, and R.H. Moss (eds), *Climate Change 1995: Impacts, Adaptations, and Mitigation of Climate Change: Scientific-Technical Analyses*, Contribution of Working Group II to the Second Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge and New York, Cambridge University Press, pp. 775-797.
- Dixon, R.K., K.J. Andrasko, F.G. Sussman, M.A. Lavinson, M.C. Trexler and T.S. Vinson. (1993). The forest-sector carbon-offset projects: near term opportunities to mitigate greenhouse gas emission. *Water, Air and Soil Pollution* 70:561-577.
- Friend, D. and Corley, R.H.V. (1994). Measuring coconut palm dry matter production. *Exp. Agric.* 30(2): 223-235.
- Frumhoff, P.C., Goetze, D.C. and Hardner, J.J. (1998). *Linking Solutions to Climate Change and Biodiversity Loss Through the Kyoto Protocol's Clean Development Mechanism*. Union of Concerned Scientists, Cambridge, MA, 14pp.
- IGBP Terrestrial Carbon Working Group. (1998). The terrestrial carbon cycle: implications for the Kyoto Protocol. *Science* 29: 1393-1394.

Lasco, R.D. and F.B. Pulhin. (2002). Forestry mitigation options in the Philippines: application of the COMAP model. ***Mitigation and Adaptation Strategies to Global Change (in press)***.

Lasco, R.D., I.Q. Guillermo, R.V.O. Cruz, N.C. Bantayan, and F.B. Pulhin. (2002). Carbon stocks assessment of a tropical secondary forest in Mt. Makiling, Philippines. *J of Tropical Forest Sci* (in press).

Lasco, R.D. and F.B. Pulhin. (2000). Forest land-use change in the Philippines and climate change mitigation. ***Mitigation and Adaptation Strategies to Global Change Journal*** 5:81-97.

Lugo, A.E. and Brown, S. (1993). Management of tropical soils as sinks and sources of atmospheric carbon. ***Plant and Soil*** 149: 27-41.

Moura-Costa P. (1996). Tropical forestry practices for carbon sequestration. In, A Zchulte and D. Schone (eds.). ***Dipterocarp Forest Ecosystems: Towards Sustainable Management***. Singapore: World Scientific. pp.308-334.

Schimmel, D., Enting, I.G., Heimann, M., Wigley, T.M.L., Rayneud, D., Alves, D. and Seigenthaler, U.: 1995, 'CO₂ and the carbon cycle', in J.T. Houghton, L.G. Meira Filho, J. Bruce, H. Lee, B.A. Callander, E. Haites, N. Harris, and K. Maskell (eds), ***Climate Change 1994: Radiative Forcing of Climate Change and an Evaluation of the IPCC IS92 Emission Scenarios***, Published for the IPCC, Cambridge University Press, Cambridge, pp. 35-71.

Trexler, M.C. and Haugen, C. (1994). ***Keeping it Green: Evaluating Tmpical Forestry Strategies to Mitigate Global Warming***, Washington D.C.: WRI, 52pp.

Watson, R.T., I.R. Noble, B. Bolin, N.H. Ravindranath, D.J. Verado and D.J. Dokken (eds) (2000). ***Land Use, Land-use Change, and Forestry***, Published for the Intergovernmental Panel for Climate Change, Cambridge University Press, Cambridge, 377pp.